

# Aboveground litterfall in Eurasian forests

LIU Chun-jiang<sup>1,2</sup>, Hannu Ilvesniemi<sup>2</sup>, Björn Berg<sup>3</sup>, Werner Kutsch<sup>1</sup>, YANG Yu-sheng<sup>4</sup>, MA Xiang-qing<sup>4</sup>, Carl J. Westman<sup>2</sup>

<sup>1</sup> Ecology Center, University of Kiel, Schauenburgerstr. 112, DE-24118 Kiel, Germany

<sup>2</sup> Department of Forest Ecology, University of Helsinki, FIN-00014 Helsinki, Finland

<sup>3</sup> Department of Soil Ecology, P.O. Box 101251, BITOEK, University of Bayreuth, DE-95 440 Bayreuth, Germany

<sup>4</sup> Department of Forestry, Fujian Agriculture and Forestry University, 353001 Nanping, Fujian, P. R. China

**Abstract:** With a data set of Eurasian forest litter fall based on 471 stands, annual litterfall was estimated to be  $6.53 \text{ Pg dm} \cdot \text{a}^{-1}$  ( $1 \text{ Pg} = 10^{15} \text{ g}$ ; dm, dry matter) in Eurasian forests, of which more than half occurred in tropical and subtropical forests, and a third in the boreal area. With litterfall, around  $2.94 \text{ Pg C}$  per year is transferred from the forest vegetation to the soil on this continent.

**Keywords:** Carbon flux; Eurasian forests; Litterfall

**CLC number:** S718.55

**Document code:** A

**Article ID:** 1007-662X(2003)01-0027-08

## Introduction

Litter fall, transferring organic matter and energy from the vegetation to the soil, is one of the major global carbon fluxes. In forest ecosystems, aboveground litter includes mainly foliage, branches, bark, and reproductive organs, and usually, foliage litterfall occupies a major fraction of total litterfall. At a continental or global scale, the synthesis of litterfall data is important for estimating and explaining quantitative litterfall, as well as its pattern.

The Eurasian continent comprises a variety of climatic conditions and forest vegetation (Ahti *et al.* 1968; Hou 1982; FAO 2001). Eurasian forests currently cover around 1.5 billion  $\text{hm}^2$ , or 41% of the total global forest area (FAO 2001) and thus play an important role in global carbon cycle. In the context of forestry management as a strategy to sequestering carbon in forest ecosystem, it is necessary to quantify the litterfall to obtain a better understanding of carbon dynamics in Eurasian forests.

Litterfall on the forest stand level has been long observed in European forests (see Rodin & Bazilevich 1967). During the period of the International Biological Program (IBP), litterfall data were collected also at some sites of Japanese and Asian tropical forests (Deangelis *et al.* 1981; Cannell 1982). Since the beginning of the 1980s, a lot of litterfall investigations have been conducted in Chinese (Zhou 1995) and Indian forests (Dadhwal *et al.* 1997). Most of these litterfall data have not yet been synthesized at a continental scale. In this synthesis, our main objectives are to estimate the annual litterfall in Eurasian forests, and to determine the

difference in litterfall between types of forests in different biomes.

## Methods

### Data Collection

To obtain original publications over litterfall in Eurasian forests, Forestry Abstract (1960s to 1998), and main forestry English-language journals (up until 1999) were searched, as well as some major reviews (Bray & Gorham 1964; Rodin & Bazilevich 1967; Cannell 1982; Proctor 1984; Berg *et al.* 1993; Dadhwal *et al.* 1997). To obtain data for China's forests, we surveyed main Chinese forestry publications from 1980 until 1998. All literature from which the data has been cited is listed in Appendix A.

In the data compilation, some criteria were followed. We used the data of forested areas (FAO 2001) to estimating litterfall in various forest groups, so the sample stands for which litterfall data was obtained in this study meet the definition of forest of FAO (2001). Stands, which had been fertilized or disturbed by forestry practice, e.g. thinning, or by e.g. fire, were not included. In the stands used, litterfall should have been collected for at least 1 year, and for the stands with the litterfall data of 2 years or more, mean values were used.

### Estimating annual litterfall

Based on an identification of ecological zones (FAO 2001), the continent's climates were classified into five domains, namely tropical, subtropical, temperate, boreal, and arctic, and the forests in each such domain were subdivided into forest types according to precipitation, temperature, and altitude. In the boreal forest the subgroup arctic forest and in tropical, subtropical and temperate areas the subgroups, steppe and desert, were excluded. Boreal forests in Europe and Asia were combined into one group, mainly due to a lower number of data in these forest

**Biography:** LIU Chun-jiang (1957-), male, research scientist, Ecology Center, University of Kiel, Schauenburgerstr. 112, DE-24118 Kiel, Germany Email: [clu@ecology.uni-kiel.de](mailto:clu@ecology.uni-kiel.de)

**Received date** 2002-11-05

**Responsible editor.** Song Funan

types. In all 13 groups of forests were used in estimating litterfall (Table 1). The area covered by each group was obtained as based on FAO's data set (FAO 2001). Correspondingly, we combined the stands from which our litterfall data were collected into 13 groups according to their geographical location, altitude, temperature, and precipitation. Mean annual total and leaf litter fall were calculated for each group and multiplied with the area of this forest group resulting in litterfall for that group on the whole continent.

**Table 1. Forest groups and the areas in estimating total litterfall in Eurasian forests.**

Forest groups	Forest area (million $\text{hm}^2$ )
<u>Tropical<sup>1</sup></u>	
Rain forest (A)	147.92
Moist (A)	36.56
Dry (A)	59.39
Mountain (A)	34.06
<u>Subtropical<sup>2</sup></u>	
Humid (A)	78.42
Dry (A)	4.00
Mountain (A)	54.42
Dry (E)	29.30
<u>Temperate<sup>3</sup></u>	
Continental (A)	45.89
Mountain (A)	37.93
Continental (E)	185.86
Oceanic (E)	28.29
<u>Boreal<sup>4</sup></u>	
Boreal (AE)	793.54
<u>Total</u>	
Total Eurasian forests	1535.57

<sup>1</sup>Not including the shrub (8.14 million  $\text{hm}^2$ ) and desert types (1.89 million  $\text{hm}^2$ ); <sup>2</sup>Not including steppe (3.23 million  $\text{hm}^2$ ) and desert (0.52 million  $\text{hm}^2$ ) types; <sup>3</sup>Not including steppe (28.79 million  $\text{hm}^2$ ) and desert (3.30 million  $\text{hm}^2$ ) types; <sup>4</sup>Not including arctic types (7.11 million  $\text{hm}^2$ ). Boreal (AE) contains coniferous, tundra and mountain types.

The letter A or E after the name of a forest group means that the group was located in Asia or Europe. The area of each forest group was based on FAO (2001).

## Results

Total litterfall ranged from  $572 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  in the Asian (A) dry forest to  $973 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  in the rain forest in the tropical area, and the proportions of leaf litter in total litterfall ranges from 53% to 73% (Table 2). Total litterfall ranged from  $438 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  in the European (E) dry forest to  $617 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  in the Mountain (A) forest with a leaf litter fraction of 70% to 79% in the subtropical area. In the temperate area, total litterfall was  $311 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  in the Asian continental forest to  $469 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  in the European continental forest and the leaf litter was 64%-87% of total litterfall. In the boreal forest, total and leaf litter fall were  $261 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  and  $188 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$ , respectively, the latter being 72% of the former.

Both the range of litterfall in a forest group and the standard deviation shows a great variation among different stands within a forest group (Table 2). For instance, in the Asian rain forest, the total litterfall ranged from  $340 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  to  $1490 \text{ g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$ , the highest thus being fourfold higher than the lowest. In the moist forest (A), the standard deviation is greater than half of the mean value (Table 2).

**Table 2. Average total litterfall and leaf litterfall in different forest groups in Eurasian forests, given as  $\text{g} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$ .**

Forest groups*	Leaf litter			Total litterfall		
	Mean±sd	Range	n	Mean±sd	Range	n
<u>Tropical</u>						
Rain forest (A) <sup>(1)</sup>	622±171	270-1075	39	973±260	340-1490	64
Moist (A) <sup>(2)</sup>	473±225	145-1070	24	714±355	300-1510	29
Dry (A) <sup>(3)</sup>	416±194	240-894	12	572±239	326-1205	15
Mountain (A) <sup>(4)</sup>	422±127	230-570	6	793±277	360-1250	17
<u>Subtropical</u>						
Humid (A) <sup>(5)</sup>	384±164	176-829	59	539±219	239-1300	64
Dry (A) <sup>(6)</sup>	380±167	84-620	21	482±267	101-1060	20
Mountain (A) <sup>(7)</sup>	443±185	167-910	17	617±208	346-1178	17
Dry (E) <sup>(8)</sup>	307±127	80-530	19	438±191	174-770	19
<u>Temperate</u>						
Continental (A) <sup>(9)</sup>	198±91	79-334	18	311±120	92-524	23
Mountain (A) <sup>(10)</sup>	312±99	219-567	10	360±129	167-667	6
Oceanic (E) <sup>(11)</sup>	301±74	125-440	28	356±95	210-671	40
Continental (E) <sup>(12)</sup>	352±108	116-520	16	469±178	233-1180	31
<u>Boreal</u>						
Boreal (E, A) <sup>(13)</sup>	188±77	23-374	83	261±108	27-508	120

Note: The letter A or E after the name of forest groups in the first column means this group is located in Asia or Europe and the number indexes the references used. The data used in this analysis were based on those of Appendix A. The numbers in the following notes refer to the papers listed in Appendix A.

- (1) Paper 14-15, 17, 34, 43-45, 63, 66, 78-79, 88, 97, 100, 118 & 136.
- (2) Paper 21, 23-24, 52, 64, 68-69, 78, 95, 100-101, 113, 128, 144-145.
- (3) Paper 2, 5, 16, 23, 42, 53 & 128.
- (4) Paper 10, 48, 51, 58, 69, 97, 128 & 146.
- (5) Paper 1, 4, 35-37, 55, 61-62, 65, 82, 84, 86-87, 91, 117, 123, 129, 132-135, 142, 144, 148 & 151.
- (6) Paper 6, 40, 42, 92, 102, 114, 119-122, 124, 130-131 & 149.
- (7) Paper 19-20, 41, 59, 93, 105-107 & 115.
- (8) Paper 10, 39-40, 51, 81 & 138.
- (9) Paper 7, 22, 27-28, 67, 116, 140-141, 143, 147 & 150.
- (10) Paper 22, 29, 49, 56, 60 & 81.
- (11) Paper 9, 25-26, 30-32, 75-76, 103 & 126-127.
- (12) Paper 8, 13, 32, 46-47, 54, 77, 108 & 125.
- (13) Paper 3, 8-9, 11, 13, 26, 32-33, 57, 70-74, 83, 89-90, 94, 99, 104, 110-112 & 138-139.

Over all Eurasia, the forested area is today more than 1 500 million  $\text{hm}^2$  (excluding arctic and desert forests), half of which was boreal forest (Table 1). Total litterfall was around 6.53 Pg dry matter per year, of which more than the half is produced in Asian tropical and subtropical forests,

and a third in the Eurasian boreal forests (Table 3). In the tropical area, most of the litterfall occurs in the rain forest ( $1.44 \text{ Pg}\cdot\text{a}^{-1}$ ), and from  $0.26 \text{ Pg}\cdot\text{a}^{-1}$  to  $0.34 \text{ Pg}\cdot\text{a}^{-1}$  is produced in the other three forest types. On the average leaf litter constituted 71% of the total litterfall in Eurasian forests.

**Table 3. Annual leaf litter and total aboveground litter production in Eurasian forests ( $10^{15} \text{ g}\cdot\text{dm}\cdot\text{a}^{-1}$ , mean  $\pm$  s.d. ).**

Forest groups	Leaf litterfall	Total litterfall
Tropical		
Rain forest (A)	$0.92\pm0.25$	$1.44\pm0.38$
Moist (A)	$0.17\pm0.08$	$0.26\pm0.13$
Dry (A)	$0.25\pm0.12$	$0.34\pm0.14$
Mountain (A)	$0.14\pm0.04$	$0.27\pm0.09$
Subtotal	1.48	2.31
Subtropical		
Humid (A)	$0.32\pm0.13$	$0.42\pm0.17$
Dry (A)	$0.02\pm0.01$	$0.02\pm0.01$
Mountain (A)	$0.24\pm0.10$	$0.34\pm0.11$
Dry (E)	$0.09\pm0.04$	$0.12\pm0.05$
Subtotal	0.67	0.94
Temperate		
Continental (A)	$0.13\pm0.06$	$0.14\pm0.06$
Mountain (A)	$0.11\pm0.01$	$0.14\pm0.05$
Oceanic (E)	$0.08\pm0.02$	$0.10\pm0.03$
Continental (E)	$0.71\pm0.13$	$0.87\pm0.33$
Subtotal	1.03	1.25
Boreal		
Boreal (AE)	$1.49\pm0.61$	$2.07\pm0.85$
Total Eurasian	4.66	6.53

## Discussion

Our estimate shows that litter production in Eurasian forests is  $6.53 \text{ Pg}$  dry mass per year, and that foliage litter occupies a major fraction (53% to 87%) (Table 2). The concentration of carbon in leaf litter ranges from 40% to 55% (e.g. Upadhyay 1990, 1993; van Wesemael 1993; Johansson 1994; Pardo *et al.* 1997; Fioretto *et al.* 1998), and this variation is not related to forest groups. To simplify the calculations, we use 0.45 as a conversion factor from litter dry matter to carbon. Thus, the carbon flux in the forest litter fall was estimated to approximate  $2.94 \text{ Pg C}$  per year.

Compared with other biomass components (e.g. stem wood, coarse branches etc.), fine litter (leaf, reproductive organs, twigs etc.) has a much higher turnover rate. Meentemeyer (1986) showed that the first-year mass-loss of fresh leaf litter ranged from 30% in boreal to 80% in tropical areas and corresponding amounts of C will be released. Today we still do not know what fractions of litter that is transferred to humus but just the mass loss on the first year is considerable. In the context of global warming, forestry management has been accepted as a strategy to mitigating atmospheric  $\text{CO}_2$  (IGBP Terrestrial Carbon Working Group 1998). In terms of carbon-fixing and

carbon-sequestering efficiencies of a forest ecosystem, litterfall and the litter turnover should be taken into account.

The global forested area was  $3.44 \times 10^9 \text{ km}^2$  in the 1990s, of which half was tropical forests ( $1.79 \times 10^9 \text{ km}^2$ ) (FAO 1995). Eurasian forests made up about  $1.50 \times 10^9 \text{ km}^2$ , being 41% of total global forests. Of the latter, boreal forests occupied a fraction of 57%, and tropical forests 26%. Global terrestrial litter production was estimated to  $39 \times 10^{15} \text{ g}\cdot\text{a}^{-1}$  (Matthews 1997) to  $54 \times 10^{15} \text{ g}\cdot\text{a}^{-1}$  (Meentemeyer *et al.* 1982), and global forest litter production was estimated to be  $29 \times 10^{15} \text{ g}\cdot\text{a}^{-1}$  (Lonsdale 1988). Our estimate showed that Eurasian forest litterfall is about one fifth of that of global forests and one ninth of that of terrestrial ecosystems. The fraction of Eurasian forest litterfall in global forest litterfall is thus much less than the fraction of Eurasian forest area of the global forest area. This may be explained by the fact that almost 60% of forests in the Eurasian continent are boreal, and that tropical forests occupy a correspondingly smaller part (27%).

We collected as complete and updated litterfall data as possible. In particular, most of the data including litterfall information in Chinese forests were published in Chinese, and thus for the first time used in a synthesis at a continental scale. For Eurasian forest litterfall, the data in our study is the most complete set published up until now. Compared with other litterfall estimates at a big geographical scale (e.g. Lonsdale 1988; Matthews 1997), the present estimate has the most abundant data set (litterfall data from 471 stands). In our analysis, we realized that more litterfall data at the stand level need to be investigated in Eurasian forests, especially in Russian boreal forests in Asia to allow more accurate estimates in the future.

## Acknowledgment

This study was financially supported by The Finnish Society of Forest Science and the Graduate School of Forest Sciences in Finland. We are grateful to X.-F. Zhou and Z.-D. Peng for helping to collect the data in Chinese publications.

## References

- Ahti, T., Hamet-Ahti, L. and Jalas, J. 1968. Vegetation zones and their sections in northwestern Europe [J]. *Ann. Bot. Fenn.*, 5: 169-211.
- Berg, B., Berg, M.P., Cortina, J., Flower-Ellis, J., Gallardo, A., Johansson, M.-B., Lundmark, J.-E. and Madeira, M. 1993. Amount of litterfall in some European coniferous forests [C]. In: Breymer, A. (Ed.) *Geography of Organic matter Production and Decay*. Warsaw: Institute of Geography and spatial Organization, Polish Academy of Sciences, p 81-110
- Bray, J.R. and Gorham, E. 1964. Litter production in forests of the world [J]. *Adv. Ecol. Res.*, 2: 101-157.
- FAO. 2001. *Forest resources assessment 2000* [M]. FAO, Rome.
- Fioretto, A. 1998. Decomposition dynamics of litters of various pine species in a Corsican pine forest [J]. *Soil Biol. Biochem.*, 30: 721-727.

- Hou Xueyu. 1982. Geography of China's Vegetation and Chemical Composition of Dominant Plants [M]. Beijing: Chinese Science Press, p 1-67. (in Chinese)
- IGBP Terrestrial Carbon Working Group. 1998. The terrestrial carbon cycle: Implications for the Kyoto Protocol [J]. *Science*, **280**: 1393-1394.
- Johansson, M.-B. 1994. Decomposition rates of Scots pine needle litter related to site properties, litter quality, and climate [J]. *Can. J. For. Res.*, **24**: 1771-1781.
- Matthews, E. 1997. Global litter production, pools, and turnover times: Estimates from measurement data and regression models [J]. *J. Geophysical Res.*, **102**(D15): 18771-18800.
- Meentemeyer, V., Box, E.O., and Thompson, R. 1982. World patterns and amounts of terrestrial plant litter production [J]. *BioScience*, **32**: 123-128.
- Meentemeyer, V. 1986. The geography of organic decomposition rates [J]. *Ann. Assoc. Amer. Geogr.*, **74**: 551-560.
- Lonsdale, W.M. 1988. Predicting the amount of litterfall in forests of the world [J]. *Ann. Botany*, **61**: 319-324.
- Pardo, F., Gil, L. and Pardos, J.A. 1997. Field study of beech (*Fagus sylvatica* L.) and melojo oak (*Quercus pyrenaica* Willd) leaf litter decomposition in the centre of the Iberian Peninsula [J]. *Plant and Soil*, **191**: 89-100.
- Proctor J. 1984. Tropical forest litterfall II. The data set [C]. In: Chadwick, A.C., Sutton, S. L. (Eds.) Tropical rain forest. Leeds: the Leeds Symposium Special Publication of the Leeds Philosophical and Literary Society, p 55-63.
- Rodin, L.E. and Bazilevich, N.I. 1967. Production and mineral cycling in terrestrial vegetation [M]. Edinburgh: Oliver and Boyd, p 20-75.
- Upadhyay, V.P. 1990. Decomposition of oak litter in a pine-mixed broadleaf forest in Himalaya [J]. *Indian Journal of Ecology*, **17**: 25-28.
- Upadhyay, V.P. 1993. Effect of initial litter quality on decomposition rates of the tree leaf litter in Himalayan forest ecosystems [J]. *Tropical Ecology*, **34**(1): 44-50.
- van Wesemael, B. 1993. Litter decomposition and nutrient distribution in humus profiles in some mediterranean forests in southern Tuscany [J]. *For. Ecol. Managem.*, **57**: 99-114.
- Zhou Xiaofeng (Ed.). 1994. Long-Term Located Research on Chinese Forest Ecosystems [C]. Harbin, Northeast Forestry University Press, p 1-10. (in Chinese)
7. Bao, C.-S. 1991. Nutrient cycling in a birch forest. In: Zhou Xiaofeng (Ed.) Located Research on the Forest Ecosystems. Harbin: Northeastern Forestry University Press, p 217-228. (in Chinese)
8. Bazilevich, N.I. 1967. Litterfall data cited in Production and mineral cycling in terrestrial vegetation (Table 9, Table 32) (Rodin L. E. and Bazilevich N. I. 1967).
9. Berg, B., Berg, M.P., Cortina, J., Flower-Ellis, J., Gallardo, A., Johansson, M.-B., Lundmark, J.-E. and Madeira, M. 1993. Amount of litterfall in some European coniferous forests [C]. In: Breymeyer, A. (Ed.) Geography of Organic Matter Production and Decay. Warsaw: Institute of Geography and Spatial Organization, Polish Academy of Sciences, p 81-110.
10. Blasco, F. and Tassy, B. 1975. Étude d'un écosystème forestier montagnard du Sud de l'Inde [J]. *Bulletin d'Ecologie*, **6**: 525-539.
11. Bonnevie-Svendsen, C. and Gjems, O. 1957. Amount and chemical composition of the litter from larch, beech, Norway spruce and Scots pine stands and its effect of the soil [J]. *Norske Skogsforsøksv.*, **48**: 111-174.
12. Bray, J.R. and Gorham, E. 1964. Litter production in forest of the world [J]. *Advances in Ecological Research*, **2**: 101-157.
13. Breymeyer, A.I. 1991. Comparative analysis of organic matter transformations in coniferous forests in Europe [C]. In: Nakagoshi, N. Golley, F.B. (Eds.) Coniferous forest ecology from an international perspective The Hague: SPB Academic Publishing bv, p 161-177.
14. Bruijnzeel, L.A. 1982. Hydrological and biogeochemical aspects of man-made forests in south-central Java, Indonesia [M]. Amsterdam: Vrije Universiteit te Amsterdam.
15. Bruijnzeel, L.A. 1985. Nutrient content of litterfall in coniferous forest plantations in central Java, Indonesia [J]. *Journal of Tropical Ecology*, **1**(4): 353-372.
16. Bunyavejchewin, S. 1989. Primary production of plots of five young closed-spaced fast-growing tree species, III. Dry matter and nutrient content of litterfall [J]. *Nat. Hist. Siam Soc.*, **37**(1). 65-73.
17. Burghouts, T., Ernsting, G., Korthals, G. and de Vris, T. 1992. Litterfall, leaf litter decomposition and litter invertebrates in primary and selectively logged dipterocarp forest in Sabah, Malaysia [J]. *Phil. Trans. R. Soc. Lon. B*, **335**: 407-416.
18. Cannell, M.G.R. 1982. World Forest Biomass and Primary Production Data [M]. London: Academic Press, p 1-150.
19. Chaturvedi, O.P. and Singh, J.S. 1987. The structure and function of pine forest in central himalaya. I. Dry matter dynamics [J]. *Annals of Botany*, **61**: 236-252.
20. Chaturvedi, O.P. and Singh, J.S. 1987. The structure and function of pine forest in central himalaya. II. Nutrient dynamics [J]. *Annals of Botany*, **61**: 253-267.
21. Chen Z.-H. 1992. Biomass and productivity of a south subtropical evergreen broadleaved forest in Heishiding Nature Conserve Area VII. Litter fall, standing crop and decomposition rate [J]. *Botanical Journal of South China*, **1**: 107-113
22. Cheng Borong. 1984. Litterfall and its nutrient content of main forest ecosystems in Changbai Mountains, Jilin [J]. *Research of Forest Ecosystem*, **4**: 19-23.
23. Crowther, J. 1987. Ecological observation in tropical karst terrain, West Malaysia. II. Rainfall interception, litterfall and nutrient cycling [J]. *Journal of Biogeography*, **14**: 145-155.
24. Dagar, J.C. and Sharma, A.K. 1993. Litterfall beneath *Bruguiera gymnorhiza* in mangrove forests of south Andamans, India [J]. *Indian Journal of Forestry*, **16**(2): 157-161.
25. de Visser, P.H.B. 1995. Effects of irrigation and balanced fertilization on nutrient cycling in a Douglas fir stand [J]. *Plant and Soil*, (168-169): 353-363.
26. de Visser, P.H.B., Beier, C., Rasmussen, L., Kreutzer, K.,

## Appendix A: Literature for Eurasian Forest Litterfall Data

1. Akai, T. and Furuno, T. 1970. Amounts of litter-fall and grazing in young loblolly pine forest [J]. *Bulletin of the Kyoto Univ. Forests* No. 42. 83-93
2. Aksornkoae, S. and Khemnark, C. 1980. Nutrient cycling in mangrove forest of Thailand [C]. In: Asian Symposium on Mangrove Environment and Management. Kuala Lumpur. University of Malaya, p 1-14.
3. Andersson, F. 1981. Biomass data. In: Reichle, D. E. (Ed.) Dynamic Properties of Forest Ecosystems [C]. Cambridge, Cambridge University Press, p 620.
4. Ando, T., Takeuchi, I., Saito, A. and Watanabe, H. 1969. Some observations of dry-matter production on the artificial two-storied stand [J]. *J. Jap. For. Soc.*, **51**: 102-107.
5. Balasubramanyan Y. 1964. Litterfall data cited in Litter Production in Forest of the World (p. 132) (Bray, J. R., and Gorham, E. 1964).
6. Bandhu, D. 1973. Chakia Project- Tropical deciduous forest ecosystem [C]. In: Kern, L. (Ed) Modeling forest ecosystems. Tennessee: Oak Ridge National Laboratory, p39-61.

- Steinberg, N., Bredemeier, M., Blanck, K., Farrell, E.P. and Cummins, T. 1994. Biological response of five forest ecosystems in the EXMAN project to input change of water, nutrients and atmospheric loads [J]. *Forest Ecology and Management*, **68**: 15-29.
27. Ding Baoyong, Song Binyan and Yao Yujun. 1986. Analysis of the litter's nutrient elements in the *Pinus koraiensis* plantation [J]. *Bulletin of Botanical Research*, **6**(4): 161-175. (in Chinese)
28. Ding Baoyong, Yao Yujun, Zhang Zongzheng and Wang Lu. 1988. Analysis of the litter's nutrient elements in the *Pinus silvestris* var. *mongolica* plantation [J]. *Bulletin of Botanical Research*, **8**(2): 143-153. (in Chinese)
29. Djalilov, K.G. and Safarov, I.S. 1981. Biomass data. In: Reichle, D.E. (Ed.) *Dynamic Properties of Forest Ecosystems*. Cambridge: Cambridge University Press.
30. Drift, J van der. 1974. Project Meerdink: Production and Decomposition of Organic Matter in an Oak Woodland [C]. Final Report 1966-71 of the Netherlands contribution to the IBP. Amsterdam: North Holland Publication Co., p 1-50.
31. Duvigneaud, P. 1970. La chenaie melangee calcicole de Virelles-Blaimont, en haute Belgique [C]. In: Duvigneaud, P. (Ed.) *Ecology and Conservation, Productivity of Forest Ecosystems* Proceedings of the Brussels Symposium, Unesco, Paris 1970, p 635-665
32. Ehwald, E. 1957. Über den Nährstoffkreislauf des Waldes [J]. *Dtsch. Akad. Landwirtschaftswissenschaften Berlin*, **6**: 1-56.
33. Finer, L. 1996. Variation in the amount and quality of litter in a *Pinus sylvestris* L. stand growing on a bog [J]. *Forest Ecology and Management*, **80**: 1-11.
34. Furtado, J.I., Verghese, S., Liew, K.S. & Lee, T.H. 1980. Litter production in a freshwater swamp forest. Tasek Bera, Malaysia [C]. In: Furtado, J. I. (Ed.) *Tropical Ecology and Development*. Proceedings of the Vth International Symposium of Tropical Ecology. Kuala Lumpur: International Society of Tropical Ecology, p 815-822
35. Furuno, T. and Saito, H. 1981. Seasonal variations of litter fall and primary consumption by herbivorous insects in *Quercus serrata* stands in Kyoto [J]. *Bulletin of the Kyoto Univ. Forests*, **53**: 52-63.
36. Furuno, T. 1993. Litter-fall and its annual and monthly fluctuations in an eastern white pine, *Pinus strobus* L., stand over twelve years [J]. *Bulletin of the Kyoto Univ. Forests*, **65**: 1-13.
37. Furuno, T. Uenishi, S. and Uenishi, K. 1979. Investigations on the productivity of Japanese fir (*Abies firma* Sieb et Zucc.) and Hemlock (*Tsuga sieboldii* Carr.) stands in Kyoto Univ. forests in Wakayama (V) Biomass and litterfall [J]. *Bulletin of the Kyoto Univ. Forests*, **51**: 58-69
38. Gallardo, J.F., Santa Regina, I., Harrison, A F. and Howard, D. M. 1995. Organic matter and nutrient dynamics in three ecosystems of the "Sierra de Bejar" mountains (Salamanca province, Spain [J]. *Acta Ecologica*, **16**(4): 447-459.
39. Garcia-Ple, C., Vanrell, P. & Morey, M. 1995. Litter fall and decomposition in a *Pinus halepensis* forest on Mallorca [J] *Journal of Vegetation Science*, **6**: 17-22.
40. Garg, R.K. and Vyas, L.N. 1975. Litter production in deciduous forest near Udaipur (South Rajasthan) India [C]. In: Golley, F.B., Medina, E. (Eds.) *Tropical Ecological Systems: Trends in Terrestrial and Aquatic Research*. New York: Springer, p 131-135.
41. Garkoti, S.C. and Singh, S.P. 1995. Variation in net primary productivity and biomass of forests in the high mountains of central Himalaya [J]. *Journal of Vegetation Science*, **6**: 23-28.
42. Gaur, J.P. and Pandey, H N. 1978. Litter production in two tropical deciduous forest communities at Varanasi, India [J]. *Oikos*, **30**(3): 570-575.
43. Gong, W.K. and Ong, J.E. 1983. Litter production and decomposition in a coastal hill dipterocarp forest [C]. In: Sutton, S., Chadwick, A. and Whitmore, T. C. (Eds.) *British Ecological Society Rain Forest Symposium*. Oxford: Blackwell Scientific Publications, p 275-285.
44. Gong, W.K. 1972. Studies on the rates of fall, decomposition and nutrient element release of leaf litter of Representative species in a lowland *Dipterocarp* forest [M]. B. Sc. Thesis, Malaya: University of Malaya, p 1-27.
45. Gong, W.K., Ong, J.E., Wong, C.H. and Dhanarajan, G. 1983. Productivity of mangrove trees and its significance in a managed mangrove ecosystems in Malaysia. In: Soepadmo, E. (Ed.) *Mangrove Environment: Research and Management*. Malaya: University of Malaya.
46. Goryshina, T.K. 1974a. Investigations of biological productivity and factors affecting it in the Les na Vorske forest-steppe oak wood [J] *Ekologija*, **3**: 5-10.
47. Goryshina, T.K. 1974b. Biological Productivity and its Factors in the Oaks of the Forest Steppe [J]. *Scientific Notes, Series of Biological Science*, No. 367.
48. Gunadi, B. 1994. Litterfall, litter turnover and respiration in two pine forest plantations in central Java, Indonesia [J]. *Journal of Tropical Forest Science*, **6**(3): 310-322.
49. Hardiwinoto, S., Yajima, T. and Igarashi, T. 1991. Stand structure and litter production of deciduous broad-leaved forests and evergreen coniferous forests in northern Hokkaido [J]. *Research Bulletins of the College Experiment Forests*, **48**(1): 115-155.
50. Hernandez, I.M., Gallardo, J.F. and Santa Regina, I. 1992. Dynamic of organic matter in forests subject to a mediterranean semi-arid climate in the Duero basin (Spain): litter production [J]. *Acta oEcologica*, **13**(1): 55-65.
51. Hladik, A. 1978. Phenology of leaf production in rain forest og Gabon: distribution and composition of food for folivores. In Montgomery, G.G. (Ed.) *The Ecology of Arboreal Folivores*. Washington D. C : Smithsonian Institute Press, p 51-71.
52. Hladik C.M. & Hladik A. 1972. Disponibilités alimentaires vitales des primates à Ceylan La Terre et la Vie, **26** 149-215
53. Hosur, G.C., Dasog, G.S. and Satyanarayana, T. 1997. Litter production and nutrient return of different tree species under plantation conditions [J] *Indian Journal of Forestry*, **20**(3): 231-235.
54. Jakucs, P. 1985. Ecology of an Oak Forest in Hungary [M] Budapest, Akademiai Kiado.
55. Kakubari, Y. 1977. Beech forests in the Nacba Mountains Distribution of primary productivity along the altitudinal gradients [J]. *JIBP Synthesis*, **16**: 201-212.
56. Kawahara, T., Sato, A., Takeuchi, I., Tadaki, Y. and Hatiya, K. 1981. Litter fall and its decomposition in a mixed stand of Japanese larch and hinoki [J]. *Bull. For & For. Prod. Res. Inst.*, (313): 79-91
57. Kazimirov, N.I. and Morozova, R.M. 1973. Biological cycling of matter in spruce forests of Karelia [M]. Leningrad. Nauka, Lenigrad Branch, Academy of Sciences, U. S S R
58. Kellman, M.C. 1970. Secondary Plant Succession in Tropical Montane Mindanao [M]. Publication BG/2, Research School of Pacific Studies. Canberra: Australia National University.
59. Khiewtam, R.S. and Ramakrishnan, P.S. 1993. Litter and fine root dynamics of a relict sacred grove forest at Cherrapunji in north-eastern India [J]. *Forest Ecology and Mangement*, **60**: 327-344.
60. Kikuzawa, K., Asai, T. and Fukuchi, M. 1984. Leaf-litter production in a plantation of *Alnus inokumae* [J]. *Journal of Ecology*, **72**: 993-999.
61. Kimura, M., Funakoshi, M., Sudo, S., Kimura, W., Yamamura, Y. and Honma, S. 1984. Litter-fall and reproductive seasonalities in a *Leucaena leucocephala* forest at Chichijima, Ogasawara (Bonin)

- Islands [J]. Bot. Mag. Tokyo, **97**: 447-455.
62. Kira, T., Ono, Y. and Hosokawa, T. 1978. Biological Production in a Warm-temperate Evergreen Oak Forest of Japan [J]. JIBP Synthesis, **18**: 201-212.
  63. Kumar, B.M. and Deepu, J.K. 1992. Litter production and decomposition dynamics in moist deciduous forests of the Western Ghats in Peninsular India [J]. Forest Ecology and Management, **50**: 181-201.
  64. Lam, P.K.S. and Dudgeon, D. 1985. Seasonal effects on litterfall in a Hong Kong mixed forest [J]. Journal of Tropical Ecology, **1**(1): 55-64.
  65. Lian, Y. and Zhang, Q. 1998. Conversion of a natural broad-leaved evergreen forest into pure and mixed plantation forests in a subtropical area: effects on nutrient cycling [J]. Can. J. For. Res., **28**: 1518-1529.
  66. Lim M.T. 1978. Litterfall and mineral nutrient content of litter in Pasoh Forest Reserve [J]. Malayan Nature Journal, **30**: 375-380.
  67. Liu Shirong, Chai Yixin, Cai Tiji and Peng Changhui. 1990. Studies on the biomass and net primary productivity of *Larix gmelini* plantations [J]. Journal of Northeastern Forestry University, **18**(2): 40-46.
  68. Lu Changyi and Lin Peng. 1990. Studies on litter fall and decomposition of *Bruguiera sexangula* (Lour.) Poir, community on Hainan Island, China [J]. Bulletin of Marine Science, **47**(1): 139-148.
  69. Lu Junpei and Liu Qihan. 1994. Litter and its decomposition process in tropical forest in Jifengling, Hainan Island [C]. In: Zhou Xiaofeng (Ed.) Long-Term Research on China's Forest Ecosystems. Harbin: Northeast Forestry University Press, p 178-191.
  70. Manakov, K.N. 1962. Characteristics of the plant litterfall in some forest plantations of the Nurmansk Province [J]. Bot. Zh., **47**(8): 30-39.
  71. Marchenko, A.I. and Karlov, Y.M. 1962. Mineral exchange in spruce forests of the Northern taiga and the forest-tundra of the Arkhangel province [J]. Pochvovedenie, (7): 45-51.
  72. Mälikonen, E. 1974. Annual primary productivity and nutrient cycle in some pine stands [J]. Communications Instituti Forestalis Fenniae (Helsinki), **84**: 7-67.
  73. Mälikonen, E. 1977. Annual primary production and nutrient cycle in a birch stand [J]. Communicationes Instituti Forestalis Fenniae (Helsinki), **91**: 5-30.
  74. Merzoev, O.G. 1981. Biomass data [C]. In: Reichle, D. E. (Ed.) Dynamic Properties of Forest Ecosystems. Cambridge: Cambridge University Press, p. 622-623.
  75. Miller, H.G., Cooper, J.M. and Miller, J.D. 1976. Effect of nitrogen supply on nutrients in litterfall and crown leaching in a stand of Corsican pine [J]. J. Appl. Ecol., **13**: 233-248.
  76. Miller, J.D., Cooper, J.M. and Miller, H.G. 1996. Amounts and nutrient weights in litterfall, and their annual cycles, from a series of fertilizer experiments on pole-stage Sitka spruce [J]. Forestry, **69**(4): 289-302.
  77. Mina, V.N. 1955. The nitrogen and ash element cycle in the oak woods of the forests-stepp [J]. Pochvovedenie, (6): 20-25.
  78. Mitchell, B.A. 1960. Comparison and assessment of site fertility under various crops and after various uses. Sub-project No. 1, deposition of organic matter [C]. Research Pamphlet 26A, p 9-11. Malaya: Forest Research Institute.
  79. Mitchell, B.A. 1964. Litterfall data cited in Litter Production in Forest of the World, p 109 (Bray & Gorham 1964).
  80. Moro, M.J., Domingo, F. and Escarré, A. 1996. Organic matter and nitrogen cycles in pine afforested catchment with a shrub layer of *Adenocarpus decorticans* and *Cistus laurifolius* in southeastern Spain [J]. Annals of Botany, **78**: 675-685.
  81. Nie, D.-P. 1985. Distribution and cycling of nutrient elements in Chinese pine plantation in Longhua region, Hebei [M]. Master thesis. Beijing: Beijing Forestry University. (in Chinese)
  82. Nie, D.-P. 1994. A comparison of productivity and nutrient cycling of Chinese fir plantations in different site conditions. J. In: Zhou Xiaofeng (Ed.) Long-Term Research on China's Forest Ecosystems. Harbin: Northeast Forestry University Press, p 123-135. (in Chinese)
  83. Nihlgård, B. 1972. Plant biomass, primary production and distribution of chemical elements in a beech and a planted spruce forest in south Sweden [J]. Oikos, **23**: 69-81.
  84. Nishira, M. 1983. Aspects of Japanese mangroves [C]. In: Proceedings of the UNESCO Regional Seminar on Human Uses of Mangrove Environment and its Management Implications. BANSDOC, Dacca, Bangladesh, p 53-59.
  85. Ogawa, H. 1978. Litterfall and mineral nutrient content of litter in Pasoh Forest [J]. Malayan Nature Journal, **30**: 375-380.
  86. Ogino, K. 1977. A beech forest at Ashiu – biomass, its increment and net production [J]. JIBP Synthesis, **16**: 201-212.
  87. Ohkubo, M., Saito, H. and Takeoka, M. 1981. Primary production in a 63-year-old plantation of *Sciadopitys verticillata* Sieb. et Zucc. at Hino, Shiga [J]. Japanese Journal of Ecology, **31**: 295-305.
  88. Ong, J.E., Gong, W.K. and Wong, C.H. 1980. Ecological Survey of the Sungai Merbok Estuarine Mangrove System [M]. School of Biological Sciences, University Sains Malaysia, Penang, Malaysia.
  89. Paavilainen, E. 1980. Effect of fertilization on plant biomass and nutrient cycle on a drained dwarf shrub pine swamp [J]. Communications Instituti Forestalis Fenniae (Helsinki), **98**: 1-23.
  90. Paavilainen, E. 1987. Effect of fertilization on the litter fall of *Pinus sylvestris* and *Betula pubescens* on drained peatlands [J]. Scand. J. For. Res., **2**: 59-75.
  91. Pande, P.K. and Sharma, S.C. 1986. Seasonality and pattern in leaf-fall and litter accretion on the forest floor in plantations of demonstration area, Forest Research Institute & College, Dehra Dun (India) [J]. Indian Forester, **112**(4): 15-18.
  92. Pandey, H. N., Gaur, J. P. and Singh, R. N. 1980. Litter input and decomposition in tropical dry deciduous forest, grassland and abandoned crop field communities at Varanasi, India [J]. Acta Oecologia Plantarum, **1**: 317-323.
  93. Pant, S.C. and Tiwari, S.C. 1992. Litter fall and litter decomposition in a montane oak forest of Garhwal Himalaya [J]. Tropical Ecology, **33**(1): 103-109.
  94. Parshchikov, A.L. 1962. The nitrogen and ash element cycle in relation to alternation of tree species in the forests of the Central Taiga [J]. Trudy Inst. Lesa Drev. Sib. Otd., **52**: 23-29 (cited in Rodin & Bazilevich 1967, Table 9).
  95. Pitman R. & Pitman J. I. 1984. Litterfall data [C]. In: Chadwick, A.C., Sutton, S.L. (Eds.) Tropical Rain Forest. Leeds: Special Publication of the Leeds Philosophical and Literary Society, p 179.
  96. Proctor, J. 1984. Tropical forest litterfall II: the data set [C]. In: Chadwick, A.C., Sutton, S.L. (Eds.) Tropical Rain Forest. Leeds: Special Publication of the Leeds Philosophical and Literary Society, p 83-116.
  97. Proctor, J., Anderson, J.M. and Vallack, H.W. 1983. Comparative studies on soils and litterfall in forests at range of altitudes on Gunung Mulu, Sarawak [J]. Malaysian Forester, **46**: 60-76.
  98. Proctor, J., Anderson, J.M., Fogden, S.C.L. and Vallack, H.W. 1983. Ecological studies in four contrasting lowland rain forests in Gunung Mulu National Park, Sarawak II. Litterfall, litter standing crop and preliminary observation on herbivory [J]. Journal of Ecology, **71**: 261-283.
  99. Pyavchenko, N.I. 1960. Biological cycle of nitrogen and ash elements in waterlogged forests [J]. Pochvovedenie No. 6. (cited in

- Rodin & Bazilevich 1967, Table 9).
100. Rai, S.N. 1981. Productivity of Tropical Rain Forests of Karnataka [M]. Ph. D. thesis, University of Bombay.
  101. Rai, S.N. and Proctor, J. 1986. Ecological studies on four rainforests in Karnaka, India [J]. Journal of Ecology, **74**: 455-463.
  102. Ranawat, M.P.S. and Vyas, L.N. 1975. Litter production in deciduous forests of Koriyat, Udipur (South Rajasthan, India [J]. Biologia (Bratislava), **30**: 41-47.
  103. Ranger, J. 1997. Nutrient dynamics during the development of a Douglas-fir (*Pseudotsuga menziesii* Mirb) stand [J]. Acta oEcologica, **18**(2): 67-90.
  104. Raulo, J. & Hokkanen, T. 1989. Litter fall of *Alnus incana* and *Alnus glutinosa* [J]. Folia Forestalia, **738**: 1-25.
  105. Rawat, Y.S., Bhatt, Y.D., Pande, P. and Singh, S.P. 1994. Production and nutrient cycling in *Arundinaria falcata* and *Lantana camara*: The two converted ecosystems in central Himalaya [J]. Tropical Ecology, **35**(1): 53-67.
  106. Rawat, Y.S. and Singh, J.S. 1988a. Structure and function of oak forest in central Himalaya. I. Dry matter dynamics [J]. Annals of Botany, **62**: 413-427.
  107. Rawat, Y.S. and Singh, J. S. 1988b. Structure and function of oak forest in central Himalaya. II. Nutrient dynamics [J]. Annals of Botany, **62**: 397-411.
  108. Remezov, N.P., Bykova, L.N. and Smirnova, K.M. 1959. Uptake and cycle of nitrogen and ash elements in the forests of European Russia. Moscow. (the data was cited in Table 9, Table 32, Production and mineral cycling in terrestrial vegetation (Rodin and Bazilevich 1967).
  109. Rodin, L.E. and Bazilevich, N.I. 1967. Production and mineral cycling in terrestrial vegetation [M]. Edinburgh: Oliver and Boyd.
  110. Rosberg, I and Stuanes, A.O. 1992. Organic matter and nutrient fluxes in Norway site. In: Johnson D.W., Linderberg S.E. (Eds.) Atmospheric Deposition and Forest Nutrient Cycling. Berlin: Springer-Verlag, p 667.
  111. Rudanova, Ye N., Tonkonogov, V.A. and Dorokhova, K.Ya. 1967. Litterfall data cited in Table 9 in Production and Mineral Cycling in Terrestrial Vegetation (see Rodin and Bazilevich 1967).
  112. Saarsalmi, A. and Mälikönen, E. 1989. Biomass production and nutrient consumption in *Alnus incana* stands [J]. Folia Forestalia, **728**: 1-16.
  113. Sasekumar, A. and Loi, J.J. 1964. Litterfall data [C]. In: Chadwick, A.C., Sutton, S.L. (Eds.) Tropical Rain Forest. Leeds: Special Publication of the Leeds Philosophical and Literary Society, p 179.
  114. Sharma, V.K. 1981. Chakia forest, Varanasi, India [C]. In: Reichle, D. E. (Ed.) Dynamic properties of Forest Ecosystems. Cambridge: Cambridge University Press, p 5-7.
  115. Sharma, E. and Ambasht, R.S. 1987. Litterfall, decomposition and nutrient release in an age sequence of *Alnus nepalensis* plantation stands in the eastern Himalaya [J]. Journal of Ecology, **75**: 997-1010.
  116. Shen Hailong, Shen Guofang and Ding Baoyong. 1996. Decomposing dynamics of several coniferous and broadleaved litters in Mongolian Scots pine plantation [J]. Scientia Silvae Sinica, **32**(5): 393-401. (in Chinese)
  117. Singh, J. and Ramakrishnan, P. S. 1982. Structure and function of a sub-tropical humid forest of Meghalaya II. Litter dynamics and nutrient cycling [J]. Proceedings of Indian Academy of Science (Plant Science), **91**: 255-268.
  118. Singh, J., George, M and Varghese, G. 1993. Nutrient cycling in moist deciduous forest-litter production and nutrient return. Indian Forester, **119**(12): 1004-1010.
  119. Singh, K.P. and Misra, R. 1978. Structure and functioning of natural, modified and silvicultural ecosystems of eastern Uttar Pradesh [R]. Technical Report. UNESCO Programme on Man and the Biosphere. Varanasi: Banaras Hindu University, India.
  120. Singh, K.P. 1968. Litter production and nutrient turnover in deciduous forests of Varanasi [C]. In Misra, R., Gopal, B. (Rds.) Proceedings of the Symposium on Recent Advances in Tropical Ecology (Part II), p 655-665.
  121. Singh, L. and Singh, J.S. 1993. Importance of short-lived components of a dry tropical forest for biomass production and nutrient cycling [J]. Journal of vegetation Science, **4**: 681-686.
  122. Singh, L. 1992. Dry matter and nutrient inputs through litter fall in a dry tropical forest of India [J]. Vegetatio **98**: 129-140.
  123. Singh, O., Sharma, D.C. and Rawat, J.K. 1993. Production and decomposition of leaf litter in sal, teak, Eucalyptus and poplar forests in Uttar Pradesh [J]. Indian Forester, **119** (2): 112-121.
  124. Singh, R.P. 1975. Biomass, nutrient and productivity structure of a stand of dry deciduous forest of Varanasi [J]. Tropical Ecology, **16**: 104-109.
  125. Smirnova, K.M. and Gorodentseva, G.A. 1958. Nutrient uptake and cycle in a birch wood [J]. Byull. Mosk. Obshch. Ispyt. Priv., **13**(12).
  126. Tavakol, M.H. and Proctor, J. 1994a. Ecological studies on three contrasting Scottish oakwoods I. Site descriptions and small litterfall [J]. Bot. J. Scotl., **47**(1): 17-32.
  127. Tavakol, M.H. and Proctor, J. 1994b. Ecological studies on three contrasting Scottish oakwoods II. Litter decomposition [J]. Bot. J. Scotl., **47**(1): 33-48.
  128. Thaiutsa, B., Suwannapinunt, W. and Kaitpraneet, W. 1978. Preliminary study of production and chemical composition of forest litter in Thailand [R]. Forest Research Bulletin 52. Bangkok: Faculty of Forestry, Kasetsart University, Thailand.
  129. Tian Dalun and Zhao Kun. 1989. Studies on the litter in a Chinese fir plantation ecosystem. I. Amount, composition and dynamics of litter [J]. Journal of Central- South Forestry College, **9**(sup.): 38-44.
  130. Tripathi, S.K. and Singh, K.P. 1994. Productivity and nutrient cycling in recently harvested and mature bamboo savannas in the dry tropics [J]. Journal of Applied Ecology, **31**: 109-124.
  131. Tripathiand, S.K. and Singh, K.P. 1996. Culm recruitment, dry matter dynamics and carbon flux in recently harvested and mature bamboo savannas in the Indian dry tropics [J]. Ecological Research, **11**: 149-164.
  132. Tsutsumi, T. 1983. On the effects of soil fertility on the rate and the nutrient element concentrations of litterfall in a forest [J]. Jap. J. Ecolo., **33**: 313-322.
  133. Tu Mengzhao. 1984. Litterfall of south-subtropical evergreen broadleaved forests in Dinghushan, Guangdong [J]. Trop. Subtrop. For. Ecosys., **1**(2): 18-21 (in Chinese)
  134. Ueda, S. and Tsutsumi, T. 1979. The amount of nutrient elements of litterfall in *Chamaecyparis obtusa* stands-influences of fertilization and site conditions [J]. Bullutin of the Kyoto Univ. Forests, **51**: 84-95.
  135. Ueda, S. and Tsutsumi, T. 1980. The amount of nutrient elements contained in litterfall of natural evergreen broadleaved forest dominated by *Machilus thunbergii* Sieb. et Zucc. [J]. Bulletin of the Kyoto Univ. Forests, **52**: 32-43.
  136. van Schaik, C.P. and Mirmanto, E. 1985. Spatial variation in the structure and litterfall of Sumatran rain forest [J]. Biotropica, **17**(3): 196-205.
  137. van Wesemael, B. and Veer, M.A.C. 1992. Soil organic matter accumulation, litter decomposition and humus forms under mediterranean-type forests in southern Tusany, Italy [J]. Journal of Soil Science, **43**: 133-144.
  138. Vedrova, E.F. 1995. Carbon pools and fluxes of 25-year old coniferous and deciduous stands in middle Siberia [C]. In: Apps, M. J.

- (Ed.) Boreal Forest and Global Change. Kluwer Academic Publishers, p 239-246.
139. Viro, P.J. 1955. Investigations on forest litter [J]. Commun. Inst. For. Fenn. **45**(6): 1-65.
  140. Wang Fengyou and Wang Yeju. 1991a. Forest litter ecology in deciduous/*Pinus koraiensis* forests. (I) Litterfall and its dynamics in the four community types. In: Zhou Xiaofeng (Ed.) Located Research on the Forest Ecosystems. Harbin: Northeast Forestry University Press, p 228-237.
  141. Wang Fengyou and Wang Yeju. 1991b. Forest litter ecology in deciduous/*Pinus koraiensis* forests: (III) Standing crops of forest litter in the four community types. In: Zhou Xiaofeng (Ed.) Located Research on the Forest Ecosystems. Harbin: Northeast Forestry University Press, p 245-249.
  142. Watanabe, H. 1978. Annual litter fall in a Nagi (*Decusscarpus nagi*) stand [J]. Bulletin of the Kyoto Univ. Forests, No. 50: 24-31.
  143. Wei Xiaohua and Zhou Xiaofeng. 1991. Nutrient cyclings in an oak forest ecosystem [C]. In: Zhou Xiaofeng (Ed.) Located Research on the Forest Ecosystems. Harbin: Northeast Forestry University Press, p 118-133. (in Chinese)
  144. Wen Yuangaung, Wei Binger and Li Jijuan. 1989. Studies on the litterfall and dynamics in some subtropical forests [J]. Scientia Silvae Sinicae, **25**(6): 542-548. (in Chinese)
  145. Wu Zhidong, Peng Fuquan, Ju Yuping, Yi Ruiling, Gu Xixian & Wu Youmei. 1990. Characteristics of the biological material cycling of several types of artificial forests and their influence on soils in south subtropical China [J]. Acta Pedologica Sinica, **27**(3): 250-261. (in Chinese)
  - Chinese)
  146. Yamada, I. 1976. Forest ecological studies of the montane forest of Mt. Pangrango, West Java III. Litter fall of the tropical forest near Cibodas [J]. South East Asian Studies, **14**: 194-229.
  147. Yao Yiantao. 1989. Studies on the biomass and nutrient cycling of the mixed plantation of *Pinus tabulaeformis* and *Platycladus orientalis* ([J]. Journal of Beijing Forestry University, **11**(3): 38-46. (in Chinese)
  148. Yu Mingjian, Chen Qichang, Li Minghong, Shi Meizhong and Cao Jianbo. 1996. Studies on the characteristics of the forest floor and the soil in the evergreen broad-leaved forest dominated by *Cyclobalanopsis glauca* in Zhejiang, China [J]. Scientia Silvae Sinicae, **32**(2): 103-110. (in Chinese)
  149. Yu Yuanchun, Yuan Honghua and Fei Shimin. 1994. Studies on litterfall of mian forest types in Xiashu forest farm, Jiangsu, China [C]. In: Zhou Xiaofeng (Ed.) Long-Term Research on China's Forest Ecosystems. Harbin: Northeast Forestry University Press, p 112-122. (in Chinese)
  150. Zhang Chenglin and Zhou Xiaofeng. 1991. Biomass of a natural secondary birch stand [C]. In: Zhou Xiaofeng (Ed.) Located Research on the Forest Ecosystems. Harbin: Northeastern Forestry University Press, p 428-435. (in Chinese)
  151. Zheng Wenjiao, Shao Cheng, Wang Liangmu and Lin Peng. 1995. Dynamic of nutrient elements in litterfall of subtropical rain forest of Hexi in Fujian [J]. Journal of Tropical and subtropical Botany, **3**(4): 38-43.